

A Second Look at Masses and Mixings of Quarks and Leptons

Berthold Stech

Institut für Theoretische Physik, Universität Heidelberg
Philosophenweg 16, D-69120 Heidelberg

Abstract

The masses of quarks and leptons in units of m_t are expressed in powers of $\sigma = \sqrt{\frac{m_c}{m_t}}$. The regularities found suggest specific mass matrices. They reproduce all masses and mixings in striking agreement with observation. A 90° relative phase angle between the off-diagonal elements of up and down quark matrices leads to a maximum magnitude for CP-violation and, for the phase angle γ of the unitary triangle, to $\gamma \simeq 90^\circ$.

Since the discovery of the top quark the masses of all quarks are known approximately. It is convenient to compare them in the $\overline{MS}$ scheme at the common scale of 1 GeV. Expressed in GeV units one has [1]

$$\begin{aligned} m_u(1 \text{ GeV}) &= (4.8 \pm 1.5)10^{-3} & m_d(1 \text{ GeV}) &= (8.5 \pm 1.5)10^{-3} \\ m_c(1 \text{ GeV}) &= 1.30 \pm 0.15 & m_s(1 \text{ GeV}) &= 0.170 \pm 0.025 \\ m_t(1 \text{ GeV}) &= 390 \pm 30 & m_b(1 \text{ GeV}) &= 6.4 \pm 0.4. \end{aligned} \quad (1)$$

This knowledge makes it possible to look for regularities of quark and lepton masses which may shed light on these mysterious quantities and on the even more mysterious mixing parameters and the CP-violating phase. One can also reconsider earlier suggestions [2]-[5] about the texture of mass matrices and their connection with the observed CP-violation.

My first observation concerns the up-quark masses. The mass ratios shown in (1) are consistent with the rule $m_u/m_c = m_c/m_t$, i. e.

$$\begin{aligned} m_t : m_c : m_u &= 1 : \sigma^2 : \sigma^4 \\ \text{with } \sigma &= \sqrt{\frac{m_c}{m_t}} = 0.058 \pm 0.004. \end{aligned} \quad (2)$$

In case there is a unique force responsible for the pattern (2) and which determines also the remaining masses and mixings, it appears natural to use σ as an expansion parameter. I take then the mass ratios (2) to hold up to $O(\sigma)$ corrections.

Expressed in powers of σ the down-quark mass ratios are consistent with the relations

$$m_b : m_s : m_d = 1 : \frac{1}{2}\sigma : 8\sigma^3 \quad (3)$$

The connection with the up-quark masses then is

$$\begin{aligned} m_b &= \alpha_{bt} \sigma m_t \\ m_s &= \alpha_{bt} \frac{1}{2} m_c = \alpha_{bt} \frac{1}{2} \sigma^2 m_t \\ m_d &= \alpha_{bt} 8 m_u = \alpha_{bt} 8 \sigma^4 m_t \\ \text{with } \alpha_{bt}(1 \text{ GeV}) &= \frac{m_b}{\sigma m_t} = 0.28 \pm 0.03. \end{aligned} \quad (4)$$

It is again suggestive to assume that the relations (3), (4) hold apart from $O(\sigma)$ corrections. Notably, in (4) the expression for m_b contains σ linearly while m_s/m_c and m_d/m_u are independent of it.

Let's then also look at the masses of the charged leptons. To a good accuracy one has

$$m_\tau : m_\mu : m_e = 1 : \sigma : \frac{3}{2}\sigma^3 \quad (5)$$

In (5) the same powers of σ appear as for the down-quark masses. Expressed in terms of m_b and m_t the lepton masses can be written

$$\begin{aligned}
m_\tau &= \alpha_{\tau b} m_b = \alpha_{\tau b} \alpha_{bt} \sigma m_t \\
m_\mu &= \alpha_{\tau b} \sigma m_b = \alpha_{\tau b} \alpha_{bt} \sigma^2 m_t \\
m_e &= \alpha_{\tau b} \frac{3}{2} \sigma^3 m_b = \alpha_{\tau b} \alpha_{bt} \frac{3}{2} \sigma^4 m_t \\
\text{with } \alpha_{\tau b}(1 \text{ GeV}) &= \frac{m_\tau}{m_b} = 0.28 \pm 0.02.
\end{aligned} \tag{6}$$

α_{bt} and $\alpha_{\tau b}$ are scale-dependent. $\alpha_{\tau b}$ approaches $\simeq 1$ at unification energies [6]. The numerical equality of $\alpha_{\tau b}$ with α_{bt} at 1 GeV (see (4) and (6)) is notable but could be entirely fortuitous.

The next task is now to construct mass matrices which appear reasonable from the point of view of the dominance of the top quark and the structures found so far.

In the standard model as well as in more general models which have no flavour-changing right-handed currents the mass matrices can be taken to be hermitian matrices without loss of generality [5], [7]. It is of advantage to take the hermitian up-quark mass matrix with eigenvalues corresponding to (2) not in its diagonal form but close to it such that the 33 element is equal to m_t up to corrections of order σ , and all other elements are of first or higher order in σ . The motivation is clear: off-diagonal matrix elements with a lower power of σ than diagonal elements and eigenvalues may have generated these by a higher order or step by step process. To express mass matrices in terms of powers of small parameters has been suggested by several authors [8]. Accordingly, the up-quark mass matrix M^U is expected to exhibit the pattern

$$M^U = \begin{pmatrix} \sim \sigma^4 & \sim \sigma^3 & \sim \sigma^2 \\ \sim \sigma^3 & \sim \sigma^2 & \sim \sigma \\ \sim \sigma^2 & \sim \sigma & 1 \end{pmatrix} m_t. \tag{7}$$

If the down-quark mass matrix M^D is indeed strongly influenced by m_t and σ as suggested by (3), (4), its 33 element to leading order will be

$$m_{33}^D = \alpha_{bt} \sigma m_{33}^U = \alpha_{bt} \sigma m_t \tag{8}$$

with all other elements being of higher order. Consequently, as a first result, all mixing angles – apart from the CP-violating phase angle – must be of order σ or smaller. In fact, the experimentally determined numerical values [9] for the Kobayashi-Maskawa matrix elements are consistent with the following expressions to be “derived” further below:

$$|V_{us}| \simeq 4\sigma, \quad |V_{cb}| \simeq \frac{\sigma}{\sqrt{2}}, \quad |V_{ub}| \simeq \sigma^2. \tag{9}$$

Numerically, (9) gives – taking $\sigma = 0.056$ –

$$|V_{us}| \simeq 0.22, \quad |V_{cb}| \simeq 0.040, \quad |V_{ub}| \simeq 0.0031 \tag{10}$$

in good agreement with present knowledge. In view of these relations it is appropriate to take σ instead of the Cabibbo angle as the expansion parameter in a Wolfenstein-type parametrization of the Kobayashi-Maskawa matrix which then reads

$$KM = \begin{pmatrix} 1 - 8A^2\sigma^2 & 4A\sigma & Be^{-i\delta}\sigma^2 \\ -4A\sigma & 1 - (8A^2 + \frac{C^2}{4})\sigma^2 & \frac{C}{\sqrt{2}}\sigma \\ (4\frac{AC}{\sqrt{2}} - Be^{i\delta})\sigma^2 & -\frac{C}{\sqrt{2}}\sigma & 1 - \frac{C^2\sigma^2}{4} \end{pmatrix} \quad (11)$$

with $A \simeq 1$, $B \simeq 1$, $C \simeq 1$. The new parametrization in terms of σ has the advantage to govern besides the KM-matrix also the quark and charged lepton mass ratios.

Let us now discuss a possible form of the hermitian up-quark mass matrix. For convenience we will take $m_{13}^U = m_{31}^{U*}$ and $m_{23}^U = m_{32}^{U*}$ to be real and positive numbers as can always be achieved by a proper choice of the quark phases. In view of the dominance of the top quark it appears natural to set

$$m_{22}^U = \pm \frac{(m_{23}^U)^2}{m_{33}}, \quad m_{12}^U = \pm \frac{m_{13}^U \cdot m_{32}^U}{m_{33}}, \quad m_{11}^U = \pm \left(\frac{m_{12}^U \cdot m_{21}^U}{m_{22}} + \frac{m_{13}^U \cdot m_{31}^U}{m_{33}} \right). \quad (12)$$

m_{22}^U , m_{12}^U and m_{11}^U are then quadratic, cubic and quartic in σ , respectively, as required by (7).

By fixing the eigenvalues of M^U (apart from signs and to leading order in σ) according to (2) the eqs. (12) have the consequence

$$\begin{aligned} m_{11}^U &= 0, \quad m_{12}^U = \pm \frac{1}{\sqrt{2}}\sigma^3 m_t, \quad m_{13}^U = \sigma^2 m_t \\ m_{22}^U &= -\frac{1}{2}\sigma^2 m_t, \quad m_{23}^U = \frac{1}{\sqrt{2}}\sigma m_t, \quad , \quad m_{33}^U = m_t. \end{aligned} \quad (13)$$

Thus, the up-quark mass matrix is, in our phase convention, a real and symmetric matrix with a rather simple structure*:

$$M^U = \begin{pmatrix} 0 & \frac{1}{\sqrt{2}}\sigma^3 & \sigma^2 \\ \frac{1}{\sqrt{2}}\sigma^3 & -\frac{1}{2}\sigma^2 & \frac{1}{\sqrt{2}}\sigma \\ \sigma^2 & \frac{1}{\sqrt{2}}\sigma & 1 \end{pmatrix} m_t. \quad (14)$$

The zero at the m_{11} position is of particular interest. It was often postulated following the classical paper by Weinberg [2]. Here it is a consequence of (12) and the negative sign of m_{22}^U .

The down-quark mass matrix M^D should not have off-diagonal elements of first order in σ as we remarked earlier. Therefore, m_{22}^D agrees to leading order in σ with

* m_{12}^U is chosen to be positive. A negative value still allowed by (12) does not give acceptable results for the KM matrix.

the corresponding eigenvalue: According to (4) and taking the sign of the eigenvalues of (14) into account, one has[†]

$$m_{22}^D = -\frac{\alpha_{bt}}{2}\sigma^2 m_t = \alpha_{bt} \cdot m_{22}^U . \quad (15)$$

With m_{33}^D of order σ and m_{22}^D given by (15) the simplest choice for the down-quark mass matrix is obtained by decoupling the b -quark from s - and d -quarks, i.e. by setting $m_{13}^D = m_{23}^D = 0$. The down-quark mass matrix then remains invariant under an arbitrary change of the b -quark phase. Taking also $m_{11}^D = 0$ to have the lightest particle mass generated by off-diagonal elements as in the case of m_u the down-quark mass matrix is given by

$$M^D = \begin{pmatrix} 0 & s_D \sigma^3 & 0 \\ s_D^* \sigma^3 & -\sigma^2/2 & 0 \\ 0 & 0 & \sigma \end{pmatrix} \alpha_{bt} m_t \quad (16)$$

with $|s_D| = 2$. The value $|s_D| = 2$ gives – to leading order in σ and apart from a removable negative sign – the desired down-quark masses (3), (4).

For the charged lepton matrix M^E we use the same texture as for M^D as suggested by models of grand unification. Then M^E is given by

$$M^E = \begin{pmatrix} 0 & s_E \sigma^3 & 0 \\ s_E^* \sigma^3 & -\sigma^2 & 0 \\ 0 & 0 & \sigma \end{pmatrix} \alpha_{\tau b} \alpha_{bt} m_t \quad (17)$$

with $|s_E| = \sqrt{\frac{3}{2}}$.

To study the CP properties resulting from (14), (16) we follow C. Jarlskog [7] and consider the commutator of the up and the down quark mass matrices and its determinant. To lowest orders in σ the commutator reads

$$[M^U, M^D] = \begin{pmatrix} \frac{s_D^* - s_D}{\sqrt{2}} \sigma^6 & \frac{1}{2}(s_D - \frac{1}{\sqrt{2}}) \sigma^5 & \sigma^3 \\ -\frac{1}{2}(s_D^* - \frac{1}{\sqrt{2}}) \sigma^5 & -\frac{s_D^* - s_D}{\sqrt{2}} \sigma^6 & \frac{1}{\sqrt{2}} \sigma^2 \\ -\sigma^3 & -\frac{1}{\sqrt{2}} \sigma^2 & 0 \end{pmatrix} \alpha_{bt} m_t^2 \quad (18)$$

and the determinant

$$Det [M^U, M^D] = \frac{1}{\sqrt{2}} (s_D^* - s_D) \sigma^{10} \alpha_{bt}^3 m_t^6 . \quad (19)$$

The determinant of the commutator measures the magnitude of CP-violation.

Because $|s_D|$ is fixed, it is immediately seen that the maximum possible amount of CP-violation will occur when the relative phase angle between the off-diagonal

[†]In ref. [5] a common proportionality factor α for the ratio of the diagonal elements of down and up quark mass matrices was postulated. Now, there is a modification for the 3rd family. As seen from (8) the ratio m_{33}^D/m_{33}^U contains the additional factor σ .

elements of the up and down quark mass matrix – in our basis the phase of s_D – will be $\pm\pi/2$:

$$s_D/|s_D| = \pm i . \quad (20)$$

As in ref. [5] and as long as experiments do not disprove it, I will stick to this attractive possibility.[‡]

The mass matrices (14), (16), (17) depend – apart from the overall scale factor m_t – on three parameters only: $\sigma = \sqrt{\frac{m_c}{m_t}}$, $\alpha_{bt} = \frac{m_b}{\sigma m_t}$ and $\alpha_{\tau b} = \frac{m_\tau}{m_b}$ with $\alpha_{bt} \simeq \alpha_{\tau b}$ at 1 GeV. But, of course, one has to keep in mind that their construction was based on mass ratios abstracted from (1). Using in addition the condition for maximal CP-violation (20), the diagonalization of the mass matrices gives 6 quark masses, 3 charged lepton masses and the 4 quark-mixing angles. To lowest order in σ one obtains, beside the mass relations we started from, the formulae already exhibited in (9) and for the phase angle γ of the triangle diagram $\gamma = \pi/2$ (by taking the minus sign in (20)).

Let us forget now the “derivation” of the mass matrices and simply propose them in the precise form given by (14), (16) (17). The special factors multiplying the powers of σ as well as the zero’s in the mass matrices are now supposed to be valid at some high energy scale. At low energies modifications occur depending on the scale difference and on the group structure between the two scales. However, apart from the scale dependence of m_t , α_{bt} and $\alpha_{\tau b}$, these modifications are generally very mild. We will ignore them here and calculate now the masses and mixings to all orders of σ by diagonalizing the mass matrices (14), (16), (17) numerically i.e. without expanding in powers of σ . Not using a best fit, but just taking the central values from (1) for m_t , α_{bt} , $\alpha_{\tau b}$ and σ , namely,

$$m_{33}^U = 390 , \quad \alpha_{bt} = \alpha_{\tau b} = 0.28 , \quad \sigma = 0.058 \quad (21)$$

one obtains the masses (in GeV units)

$$\begin{aligned} m_t(1 \text{ GeV}) &= 391 & m_c(1 \text{ GeV}) &= 1.31 & m_u(1 \text{ GeV}) &= 4.4 \cdot 10^{-3} \\ m_b(1 \text{ GeV}) &= 6.33 & m_s(1 \text{ GeV}) &= 0.19 & m_d(1 \text{ GeV}) &= 9.4 \cdot 10^{-3} \\ m_\tau(1 \text{ GeV}) &= 1.77 & m_\mu(1 \text{ GeV}) &= 0.103 & m_e(1 \text{ GeV}) &= 0.52 \cdot 10^{-3} \end{aligned} \quad (22)$$

and for the Kobayashi-Maskawa matrix elements and the angles α , β , γ of the unitarity triangle

$$\begin{aligned} |V_{us}| &= 0.216 , & |V_{cb}| &= 0.041 , & |V_{ub}| &= 0.0034 , & |V_{td}| &= 0.0094 , \\ \alpha &= 70^\circ , & \beta &= 20^\circ , & \gamma &= 90^\circ . \end{aligned} \quad (23)$$

[‡]In ref. [5] it was suggested that there exists a current quark basis in which all off-diagonal elements of the down-quark mass matrix are purely imaginary (using a diagonal up-quark mass matrix). To lowest order in σ the requirement (20) is fully equivalent to this earlier proposal, as can be seen by transforming M^U and M^D such that M^U is diagonal and by using appropriate quark phases.

In conventional notation of the KM matrix V_{ub} and V_{td} read

$$V_{ub} = -0.0034i, \quad V_{td} = -0.0033i + 0.0088. \quad (24)$$

The agreement with known data [9] is quite encouraging.

I can say little about lepton (neutrino) mixing. It depends of course on the neutrino mass matrix for which we have no good information at present. With the charged lepton matrix in the form (16) the neutrino mass matrix depending on m_t and σ will certainly be close to a diagonal matrix. Thus, only small mixing angles, the largest definitely smaller than the Cabibbo angle, can be expected. The similarity of M^E with M^D suggests CP-violation and $s_E/|s_E| = \pm i$ if maximal CP-violation occurs in the quark sector.

In conclusion I can state: The masses and mixings of quarks and leptons appear to have a common origin which suggests mass matrices of a particularly simple form. The elements of the mass matrices are governed by powers of the small parameter $\sigma = \sqrt{\frac{m_c}{m_t}}$, which is close to $\frac{2m_s}{m_b}$ and $\frac{m_\mu}{m_\tau}$. The factors in front of the powers of σ remind of Clebsch-Gordan or normalization-type numbers. Their knowledge may possibly help to find the underlying dynamics. For instance, one might interpret these factors as manifestations of a family-type symmetry in a similar way as the numbers obtained from models studied in ref. [10].

Acknowledgement: It is a pleasure to thank D. Gromes, M. Jamin, O. Nachtmann, M.G. Schmidt and C. Wetterich for valuable remarks.

References

- [1] M. Jamin, private communication
- [2] S. Weinberg, Ann. N. Y. Academy of Sc. **38** (1977) 1945
- [3] H. Georgi and D. V. Nanopoulos, Nucl. Phys. **B155** (1979) 52
- [4] H. Fritzsch, Nucl. Phys. **B155** (1979) 189
- [5] B. Stech, Phys. Lett. **130B** (1983) 189
- [6] M. S. Chanowitz, J. Ellis, and M. K. Gaillard, Nucl. Phys. **B128** (1977) 506;
A. Buras, J. Ellis, M. K. Gaillard, and D. V. Nanopoulos, Nucl. Phys. **B135** (1978) 66
- [7] C. Jarlskog, Phys. Rev. Lett. **15**, (1985) 1039 and Z. f. Phys. **C29** (1985) 491
- [8] C. D. Froggatt and H. B. Nielsen, Nucl. Phys. **B147** (1979) 277;
S. Dimopoulos, Phys. Lett. **B129** (1983) 417;

- J. Bijnens and C. Wetterich, Nucl. Phys. **B283** (1987) 237, Phys. Lett. **B199** (1987) 525;
 L. E. Ibanez and G. G. Ross, Phys. Lett. **B332** (1994) 100
- [9] Particle Data group and T. E. Browder, K. Honscheid, and D. Pedrini, OHSTPY-HEP-E-96-006
- [10] L. E. Ibanez and G. G. Ross, Phys. Lett. **B332** (1994) 100;
 M. Carena, S. Dimopoulos, C.E. M. Wagner, and S. Raby, Phys. Rev. **D52** (1995) 4133;
 C. D. Froggatt, hep-ph 9603432;
 C. D. Froggatt, H. B. Nielsen, and D. J. Smith, Phys. Lett. **B385** (1996) 150